

EARLY START FOR BETTER¹

POSSIBILITIES

Legacy effect²

This is a model and not a patient

Legacy effect underscores the importance of **Early Initiation** of intensive treatment in people with **Type 2 Diabetes**.²



Early Initiation of oral semaglutide provides both robust **glycaemic control** and **weight loss**.³



Oral semaglutide provides both **weight loss** of up to **5 kg**, **HbA1c reduction** of up to **1.5%**.^{4,5}



Legacy effect may help prevent long term diabetes complications.²



Visit HCP Portal to learn more

Abbreviations:

T2DM: Type 2 diabetes Mellitus,
HbA1c: Glycated hemoglobin

References:

1. ADA, Standards of Care in Diabetes-2024. Diabetes Care 1 January 2024; 47
2. Holman RR, et al. N Engl J Med. 2008 Oct 9;359(15):1577-89.
3. Goldenberg R et al. Can J Diabetes. 2021; 45(7):S28-29
4. Aroda VR et al. Diabetes Care. 2019 Sep;42(9):1724-1732.
5. Pratley R et al. Lancet. 2019;394:39-50 16.

Abbreviated prescribing information

[Click here to read Rybelsus API](#)

Disclaimer:

Apis bull is a registered trademark of Novo Nordisk A/S.
Not real images of patients, only for illustrative purposes.

For the use only of registered medical practitioner
or a hospital or a laboratory